

Tylenchorhynchus claytoni Steiner 1937

THE TOBACCO STUNT NEMATODE

J. B. MacGowan

Tylenchorhynchus claytoni, the tobacco stunt nematode, was described in 1937 by G. Steiner from tobacco root specimens which were collected by E. E. Clayton in Florence, South Carolina (16, 23). This nematode is a migratory ectoparasite and is known to feed upon the roots of a wide variety of plant hosts (15). It is hosted by or reported to be associated with at least 100 species of plants (2, 13, 14, 15, 16, 17, 18, 23, 26). A host list is being prepared for publication at a later date.

In the United States the tobacco stunt nematode has been reported from Alabama, Arkansas, California, Colorado, Connecticut, Florida, Georgia, Kentucky, Louisiana, Maryland, Massachusetts, Mississippi, North Carolina, Oregon, Rhode Island, Texas, Virginia, West Virginia, and Wisconsin (1, 2, 5, 8, 9, 16, 18, 25, 27). It has also been reported from Belgium, Canada, Denmark, Great Britain, Holland, India, Japan, Netherlands, and Switzerland (1, 16, 20).

Tylenchorhynchus claytoni, as a parasite of tobacco, can cause moderate stunting with reduction of top growth (7). Investigations indicate that roots of infected tobacco plants are stunted and shriveled and display loss of turgor. They appear to grow in a dense clumped mass and evidence neither observable lesions nor decay (10). In varieties of tobacco susceptible to wilt caused by Fusarium oxysporum (Schlecht.) Wr. var. nicotianae J. Johnson, the tobacco stunt nematode is known to increase the incidence of wilt (11, 20, 26).

This nematode has been demonstrated to be pathogenic to some grasses and is associated with nematode damage to others. Where pathogenicity has been proven, symptoms of root injury have ranged from destruction below a depth of 10 cm, to no observable root injury (9, 13, 15, 16, 19).

The tobacco stunt nematode is a common parasite of azaleas. Nematode activity is reflected in smaller root systems, reduced top growth, and, in some instances, chlorosis (3, 4, 5, 22). Further, this nematode can predispose azaleas to root infection and rot by Phytophthora cinnamomi Rands (6). It is parasitic to a variety of pine species and is associated with unthriftiness, chlorosis, and death of pine nursery seedlings (12, 16, 19, 20, 21, 24). It can lead to reduced root systems in corn and has been associated with stunting in the field (15, 18).

Survey and Detection:

1. Examine top parts of plants for stunting, chlorosis, or general unthriftiness.
2. Submit approximately one pint of soil from the root area of the plant to a nematology laboratory.

Literature Cited

1. Allen, M. W. 1955. A review of the nematode genus Tylenchorhynchus. Univ. of Calif. Publications in Zoology 61:129-166.
2. Anonymous. 1960. Distribution of plant parasitic nematodes in the south. South. Coop. Series Bull. 74 (Agric. Exp. Sta. of Alabama and other Southern States). 72p.
3. Barker, K. R., and G. Worf. 1964. Parasitism of 'southern stock' azaleas in Wisconsin by Tylenchorhynchus claytoni, Trichodorus christiei, and Meloidogyne incognita. Phytopathology 54:887. (Abstr.).
4. _____. 1966. Effect of nutrients on nematode activity on azalea. Phytopathology 56:1024-1027.
5. _____, and A. H. Epstein. 1965. Nematodes associated with decline of azaleas in Wisconsin. Plant Dis. Rept. 49:47-49.

6. Daykin, M. E., and D. M. Benson. 1977. Interaction of Phytophthora cinnamomi and Tylenchorhynchus claytoni in azalea decline. Proc. Amer. Phytopath. Soc. 4:126. (Abstr.).
7. Deubert, K. H., R. L. Norgren, S. M. Paracer, and B. M. Zuckerman. 1967. The influence of Tylenchus agricola and Tylenchorhynchus claytoni on corn roots under gnotobiotic conditions. Nematologica 13:56-62.
8. Esser, R. P. 1967. Foliar and other plant-parasitic nematodes associated with azalea in Florida. Plant Dis. Rept. 51:46-48.
9. Feldmesser, J., and A. M. Golden. 1972. Control of nematodes damaging home lawn grasses in two counties in Maryland. Plant Dis. Rept. 56:476-480.
10. Graham, T. W. 1954. The tobacco stunt nematodes in South Carolina. Phytopathology 44:332. (Abstr.).
11. Holdeman, Q. L. 1956. The effect of the tobacco stunt nematode on the incidence of Fusarium wilt in flu-cured tobacco. Phytopathology 46:129. (Abstr.).
12. Hopper, B. E. 1958. Plant parasitic nematodes in the soils of southern forest nurseries. Plant Dis. Rept. 42:308-314.
13. Hoveland, C. S., R. Rodriguez-Kabana, and C. D. Berry. 1975. Phalaris and tall fescue forage production as affected by nematodes in the field. Agronomy J. U.S.A. 67:714-717.
14. Khera, S., and B. M. Zuckerman. 1963. In vitro studies of host parasite studies of some plant parasitic nematodes. Nematologica 9:1-6.
15. Krusberg, L. R. 1959. Investigations on the life cycle, reproduction, feeding habits, and host range of Tylenchorhynchus claytoni Steiner. Nematologica 4:187-197.
16. Loof, P. A. A. 1974. Tylenchorhynchus claytoni. C. I. H. descriptions of plant parasitic nematodes. Set 3, No. 39. Commonwealth Agricultural Bureaux.
17. Milne, D. L. 1972. Nematodes of tobacco, pp. 159-186. In Economic Nematology. J. M. Webster (ed.), Academic Press. New York. 553p.
18. Nelson, R. R. 1955. Nematode parasites of corn in the coastal plain of North Carolina. Plant Dis. Rept. 39:818-819.
19. Reuhl, J. L. 1966. Nematodes parasitic on forest trees I. Reproduction of ectoparasites on pines. Nematologica 12:443-447.
20. _____. 1972. Nematodes of forest trees, pp. 312-334. In Economic Nematology. J. M. Webster (ed.), Academic Press. New York. 563p.
21. _____. 1973. Influence of plant parasitic nematodes on pine seedlings. J. Nematol. 5:7-9.
22. Sher, S. A. 1958. The effect of nematodes on azaleas. Plant Dis. Rept. 42:84-85.
23. Steiner, F. 1937. Opuscula miscellanea nematologica, V. Proc. Helminth. Soc. Wash. 4:33-34.
24. Sutherland, J. R., and R. E. Adams. 1965. The parasitism of red pine and other forest nursery crops by Tylenchorhynchus claytoni Steiner. Nematologica 10:637-643.
25. Tarjan, A. C. 1953. Known and suspected plant parasitic nematodes of Rhode Island. Proc. Helminth. Soc. Wash. 20:49-54.
26. Thorne, G. 1961. Principles of nematology. McGraw-Hill. New York. 553p.
27. Troll, J. R., and R. A. Rohde. 1966. Pathogenicity of Pratylenchus penetrans and Tylenchorhynchus claytoni on turfgrasses. Phytopathology 56:995-998.